

SHEP 92/93-18

hep-ph/9305282

Squark Contributions to Higgs Boson Masses in the Next-to-Minimal Supersymmetric Standard Model

T. Elliott, S. F. King and P. L. White

Physics Department

University of Southampton

Southampton

SO9 5NH

U.K.

Abstract

Within the context of an effective potential formalism we calculate the contribution to Higgs boson masses in the next-to-minimal supersymmetric standard model from squark loops. We then supplement a previously performed renormalisation group analysis of the Higgs sector of this model with these results in order to determine the shift in the bound on the lightest CP-even Higgs boson mass as a result of squark effects. The improved bound on the lightest neutral CP-even Higgs boson mass, including squark contributions, is $m_h \leq 146$ (139, 149) GeV for $m_t = 90$ (140, 190) GeV. For $m_t = 190$ GeV squark effects contribute 23 GeV to the bound, with smaller contributions for small m_t values.

The next-to-minimal supersymmetric standard model [1, 2] (NMSSM) provides an elegant extension of the minimal supersymmetric standard model [3] (MSSM) which eliminates the μ -problem, the appearance of a mass-scale in the superpotential. The NMSSM differs from the MSSM by the presence of a gauge singlet field whose vacuum expectation value (vev) plays the role of the mass parameter μ in the MSSM, and is defined by the superpotential

$$W = h_t Q H_2 t^c + \lambda N H_1 H_2 - \frac{1}{3} k N^3 + \dots, \quad (1)$$

where the superfield $Q^T = (t_L, b_L)$ contains the left-handed top and bottom quarks, and t^c contains the charge conjugate of the right-handed top quark; H_1 and H_2 are the usual Higgs doublet superfields and N is the Higgs gauge singlet superfield; the ellipsis represents terms whose relatively small couplings play no role in our analysis (in particular, we assume that the bottom quark Yukawa coupling, h_b , satisfies $h_b \ll h_t$). The fields $H_1^T = (H_1^0, H_1^-)$, $H_2^T = (H_2^+, H_2^0)$ and N develop vevs which may be assumed to be of the form [1]

$$\langle H_1 \rangle = \begin{pmatrix} \nu_1 \\ 0 \end{pmatrix}, \quad \langle H_2 \rangle = \begin{pmatrix} 0 \\ \nu_2 \end{pmatrix}, \quad \langle N \rangle = x, \quad (2)$$

where ν_1 , ν_2 and x are real, and $\sqrt{\nu_1^2 + \nu_2^2} = \nu = 174$ GeV. The low energy physical spectrum of the Higgs scalars consists of 3 CP-even neutral states, 2 CP-odd neutral states, and 2 charged scalars. A third CP-odd state is a Goldstone mode which becomes the longitudinal component of the Z^0 , while a further two charged scalars become those of the $W^\pm$'s.

The tree-level bound on the lightest CP-even Higgs scalar mass in the NMSSM [2],

$$m_h^2 \leq M_Z^2 + (\lambda^2 \nu^2 - M_Z^2) \sin^2 2\beta, \quad (3)$$

where $\tan\beta = \nu_2/\nu_1$, and radiative corrections to it [4, 5], have been the subject of much discussion. In Ref. [4] renormalisation group (RG) analyses were performed assuming either supersymmetric (SUSY) RG equations down to near the top quark mass, or the existence of only one light Higgs boson with a mass below the SUSY breaking scale. These analyses assumed degenerate squarks decoupling below their common mass scale M_{SUSY} . In Refs. [5, 6], within the effective potential formalism, squark effects have been considered in the NMSSM. However, no systematic attempt was made to study their effects on the bound in Eq. (3), but rather one slice through parameter space was examined [6]. Moreover, radiative corrections to the charged scalar mass-squared matrix were not presented, though these do not change the bound.

Recently we performed a low energy renormalisation group analysis of the Higgs sector of the NMSSM and found that $m_h \leq 146$ GeV [7]. This analysis considered the more general case of all the Higgs bosons having masses below the SUSY breaking scale (M_{SUSY}), and used non-SUSY RG equations below M_{SUSY} . However, in that analysis the squarks were assumed to be degenerate at $M_{SUSY} = 1$ TeV. Here we supplement this calculation with a computation of the stop and, where necessary, sbottom contributions to the CP-even, CP-odd and charged mass-squared matrices, from which we may determine modifications to the bound due to a general spectrum of squarks. We then attempt systematically to examine the squark corrected bound by surveying as much of the parameter space as practicable.

Our previous analysis worked exclusively with $V_0(\mu)$, $\mu = 150$ GeV being approximately the bound on the lightest CP-even state. From the superpotential in Eq. (1) we may obtain the pure Higgs part of V_0 , where we have renormalised at the $\overline{MS}$ scale μ :

$$\begin{aligned}
V_0(\mu) &= \frac{1}{2}\lambda_1(H_1^\dagger H_1)^2 + \frac{1}{2}\lambda_2(H_2^\dagger H_2)^2 + (\lambda_3 + \lambda_4)(H_1^\dagger H_1)(H_2^\dagger H_2) \\
&- \lambda_4|H_2^\dagger H_1|^2 + \lambda_5|N|^2|H_1|^2 + \lambda_6|N|^2|H_2|^2 \\
&+ \lambda_7(N^{*2}H_1H_2 + h.c.) + \lambda_8|N|^4 \\
&+ m_1^2|H_1|^2 + m_2^2|H_2|^2 + m_3^2|N|^2 \\
&- m_4(H_1H_2N + h.c.) - \frac{1}{3}m_5(N^3 + h.c.),
\end{aligned} \tag{4}$$

where $\lambda_i = \lambda_i(\mu)$ and $m_i = m_i(\mu)$. At M_{SUSY} the couplings λ_i and m_i are given by:

$$\begin{aligned}
\lambda_1 &= \lambda_2 = (g_2^2 + g_1^2)/4, \quad \lambda_3 = (g_2^2 - g_1^2)/4, \\
\lambda_4 &= \lambda^2 - g_2^2/2, \quad \lambda_5 = \lambda_6 = \lambda^2, \quad \lambda_7 = -\lambda k, \quad \lambda_8 = k^2, \\
m_1 &= m_{H_1}, \quad m_2 = m_{H_2}, \quad m_3 = m_N, \quad m_4 = \lambda A_\lambda, \quad m_5 = k A_k.
\end{aligned} \tag{5}$$

g_1 and g_2 are the $U(1)_Y$ and $SU(2)_L$ gauge coupling constants, respectively, and m_{H_i} , m_N , A_λ and A_k are soft SUSY breaking parameters. We use the RG equations [7] to allow to us calculate the couplings renormalised at $\mu = 150$ GeV given their values at M_{SUSY} .

In order to calculate the shifts in the mass-squared matrices due to squarks, we employ the full one-loop effective potential. Much of our analysis of the squark spectrum is similar to that of the MSSM in Ref. [8], whose notation we follow closely. The one-loop effective potential is given by

$$\begin{aligned}
V_1(Q) &= V_0(Q) + \Delta V_1(Q), \\
\Delta V_1(Q) &= \frac{1}{64\pi^2} \text{Str} \mathcal{M}^4 \left(\ln \frac{\mathcal{M}^2}{Q^2} - \frac{3}{2} \right),
\end{aligned} \tag{6}$$

where $V_0(Q)$ is the tree-level potential at the arbitrary $\overline{MS}$ scale Q , Str denotes the usual supertrace, and $\mathcal{M}^2$ is the field-dependent mass-squared matrix. We shall

take $Q = M_{SUSY} = 1$ TeV.¹ V_0 is restricted to the pure Higgs part of the tree-level potential. Since our RG approach has already included the logarithmic effects of top quark and Higgs boson loops we exclude these particles from the supertrace. The logarithmic contributions to $\Delta V_1(Q)$ from the top quark and Higgs bosons may be absorbed into $V_0(Q)$ where $Q = 1$ TeV, to yield $V_0(\mu)$ where $\mu = 150$ GeV, as discussed above. It only remains to calculate $\Delta V_1(Q)$, where $Q = 1$ TeV, involving squark contributions.

To calculate the shifts in the Higgs mass matrices as a result of squark effects we require the field-dependent mass-squared matrices of the squarks. Taking the contribution to the soft SUSY breaking potential involving squarks to be

$$\Delta V_{soft} = h_t A_t (\tilde{Q} H_2 \tilde{t}^c + h.c.) + m_Q^2 |\tilde{Q}|^2 + m_T^2 |\tilde{t}^c|^2 + m_B^2 |\tilde{b}^c|^2, \quad (7)$$

where b^c contains the charge conjugate of the right-handed bottom quark, and tildes denote the scalar components of the superfields, together with the superpotential in Eq. (1), leads to the field-dependent squark mass-squared matrix (ignoring contributions proportional to gauge couplings and h_b)

$$\mathcal{M}^2 = \begin{pmatrix} m_Q^2 + h_t^2 |H_2^0|^2 & \lambda h_t N H_1^0 + h_t A_t \bar{H}_2^0 & -h_t^2 \bar{H}_2^0 H_2^+ & 0 \\ \lambda h_t \bar{N} \bar{H}_1^0 + h_t A_t H_2^0 & m_T^2 + h_t^2 (|H_2^0|^2 + H_2^+ H_2^-) & \lambda h_t \bar{N} H_1^+ - h_t A_t H_2^+ & 0 \\ -h_t^2 H_2^0 H_2^- & \lambda h_t N H_1^- - h_t A_t H_2^- & m_Q^2 + h_t^2 H_2^+ H_2^- & 0 \\ 0 & 0 & 0 & m_B^2 \end{pmatrix}, \quad (8)$$

in the basis $\{\tilde{t}_L, \tilde{t}_R^c, \tilde{b}_L, \tilde{b}_R^c\}$, where a bar denotes complex conjugation. Manifestly, one eigenvalue is field-independent and may be discarded. Only the upper 2×2 submatrix contributes to the CP-even and CP-odd mass-squared matrices, since the charged fields have zero vevs in order not to break QED, whereas the upper 3×3 submatrix contributes to the charged mass-squared matrix. By differentiating ΔV_1 once with respect to the fields we may obtain the shifts in the minimisation conditions

¹This value of Q is of course totally arbitrary. The value of 1 TeV is chosen to be consistent with our previous analysis. The results are independent of Q to one loop order.

induced by squark loops, and twice will yield the shifts in the mass matrices.²

In the basis $\{H_1, H_2, N\}$ we have the following mass-squared matrices, after ensuring that the full one-loop potential is correctly minimised. The couplings λ_i are those obtained from the potential in Eq. (4) renormalised at the scale $\mu = 150$ GeV. For notational simplicity we drop the tildes. Moreover, we work in the basis of mass eigenstates $\{t_1, t_2, b_1, b_2\}$ (though, since h_b , the bottom quark Yukawa coupling, is taken to be zero, b_2 never contributes to the mass-squared matrices), where the mass of the t_1 squark is m_{t_1} , and so on for the other squarks. The CP-even (scalar) mass-squared matrix is

$$M_s^2 = M^2 + \delta M^2, \quad (9)$$

where

$$M^2 = \begin{pmatrix} 2\lambda_1\nu_1^2 & 2(\lambda_3 + \lambda_4)\nu_1\nu_2 & 2\lambda_5x\nu_1 \\ 2(\lambda_3 + \lambda_4)\nu_1\nu_2 & 2\lambda_2\nu_2^2 & 2\lambda_6x\nu_2 \\ 2\lambda_5x\nu_1 & 2\lambda_6x\nu_2 & 4\lambda_8x^2 - m_5x \end{pmatrix} + \begin{pmatrix} \tan\beta[m_4x - \lambda_7x^2] & -[m_4x - \lambda_7x^2] & -\frac{\nu_2}{x}[m_4x - 2\lambda_7x^2] \\ -[m_4x - \lambda_7x^2] & \cot\beta[m_4x - \lambda_7x^2] & -\frac{\nu_1}{x}[m_4x - 2\lambda_7x^2] \\ -\frac{\nu_2}{x}[m_4x - 2\lambda_7x^2] & -\frac{\nu_1}{x}[m_4x - 2\lambda_7x^2] & \frac{\nu_1\nu_2}{x^2}[m_4x] \end{pmatrix} \quad (10)$$

and

$$\delta M^2 = \begin{pmatrix} \Delta_{11}^2 & \Delta_{12}^2 & \Delta_{13}^2 \\ \Delta_{12}^2 & \Delta_{22}^2 & \Delta_{23}^2 \\ \Delta_{13}^2 & \Delta_{23}^2 & \Delta_{33}^2 \end{pmatrix} + \begin{pmatrix} \tan\beta & -1 & -\frac{\nu_2}{x} \\ -1 & \cot\beta & -\frac{\nu_1}{x} \\ -\frac{\nu_2}{x} & -\frac{\nu_1}{x} & \frac{\nu_1\nu_2}{x^2} \end{pmatrix} \Delta_p^2. \quad (11)$$

The Δ_{ij}^2 and Δ_p^2 are given by

$$\begin{aligned} \Delta_p^2 &= \frac{3}{16\pi^2} h_t^2 (\lambda x) A_t f(m_{t_1}^2, m_{t_2}^2), \\ \Delta_{11}^2 &= \frac{3}{8\pi^2} h_t^4 \nu_2^2 (\lambda x)^2 \cdot \left(\frac{A_t + \lambda x \cot\beta}{m_{t_2}^2 - m_{t_1}^2} \right)^2 g(m_{t_1}^2, m_{t_2}^2), \\ \Delta_{22}^2 &= \frac{3}{8\pi^2} h_t^4 \nu_2^2 \cdot \left(\ln \frac{m_{t_1}^2 m_{t_2}^2}{M_{SUSY}^2} + \frac{2A_t(A_t + \lambda x \cot\beta)}{m_{t_2}^2 - m_{t_1}^2} \ln \frac{m_{t_2}^2}{m_{t_1}^2} \right) \\ &\quad + \frac{3}{8\pi^2} h_t^4 \nu_2^2 \cdot \left(\frac{A_t(A_t + \lambda x \cot\beta)}{m_{t_2}^2 - m_{t_1}^2} \right)^2 g(m_{t_1}^2, m_{t_2}^2), \\ \Delta_{33}^2 &= \frac{3}{8\pi^2} h_t^4 \nu_2^2 (\lambda \nu_1)^2 \cdot \left(\frac{A_t + \lambda x \cot\beta}{m_{t_2}^2 - m_{t_1}^2} \right)^2 g(m_{t_1}^2, m_{t_2}^2), \end{aligned}$$

²In fact, this is only approximately true due to Higgs self-energy corrections; these are expected to be small for the lightest Higgs bosons.

$$\begin{aligned}
\Delta_{12}^2 &= \frac{3}{8\pi^2} h_t^4 \nu_2^2 (\lambda x) \cdot \left(\frac{A_t + \lambda x \cot \beta}{m_{t_2}^2 - m_{t_1}^2} \right) \left(\ln \frac{m_{t_2}^2}{m_{t_1}^2} + \frac{A_t(A_t + \lambda x \cot \beta)}{m_{t_2}^2 - m_{t_1}^2} g(m_{t_1}^2, m_{t_2}^2) \right), \\
\Delta_{13}^2 &= \frac{3}{8\pi^2} h_t^4 \nu_2^2 (\lambda x) \cdot (\lambda \nu_1) \cdot \left(\frac{A_t + \lambda x \cot \beta}{m_{t_2}^2 - m_{t_1}^2} \right)^2 g(m_{t_1}^2, m_{t_2}^2) \\
&\quad - \frac{3}{8\pi^2} h_t^2 (\lambda x) \cdot (\lambda \nu_1) \cdot f(m_{t_1}^2, m_{t_2}^2), \\
\Delta_{23}^2 &= \frac{\nu_1}{x} \Delta_{12}^2,
\end{aligned} \tag{12}$$

where the functions f and g are defined by

$$\begin{aligned}
f(m_{t_1}^2, m_{t_2}^2) &= \frac{1}{m_{t_2}^2 - m_{t_1}^2} \left[m_{t_1}^2 \ln \frac{m_{t_1}^2}{M_{SUSY}^2} - m_{t_1}^2 - m_{t_2}^2 \ln \frac{m_{t_2}^2}{M_{SUSY}^2} + m_{t_2}^2 \right], \\
g(m_{t_1}^2, m_{t_2}^2) &= \frac{1}{m_{t_1}^2 - m_{t_2}^2} \left[(m_{t_1}^2 + m_{t_2}^2) \ln \frac{m_{t_2}^2}{m_{t_1}^2} + 2(m_{t_1}^2 - m_{t_2}^2) \right].
\end{aligned} \tag{13}$$

If the squarks are degenerate in mass, then equation Eq. (8) implies that $A_t + \lambda x \cot \beta = 0$. Furthermore, if $m_{t_1} = m_{t_2} = M_{SUSY}$, then Eq. (11) and Eq. (12) imply that the squark contribution to the CP-even mass-squared matrix vanishes. This is the limit of our previous analysis.

The CP-odd (pseudoscalar) mass-squared matrix is

$$M_p^2 = \tilde{M}^2 + \delta \tilde{M}^2, \tag{14}$$

where

$$\tilde{M}^2 = \begin{pmatrix} \tan \beta [m_4 x - \lambda_7 x^2] & [m_4 x - \lambda_7 x^2] & \frac{\nu_2}{x} [m_4 x + 2\lambda_7 x^2] \\ [m_4 x - \lambda_7 x^2] & \cot \beta [m_4 x - \lambda_7 x^2] & \frac{\nu_1}{x} [m_4 x + 2\lambda_7 x^2] \\ \frac{\nu_2}{x} [m_4 x + 2\lambda_7 x^2] & \frac{\nu_1}{x} [m_4 x + 2\lambda_7 x^2] & 3m_5 x + \frac{\nu_1 \nu_2}{x^2} [m_4 x - 4\lambda_7 x^2] \end{pmatrix} \tag{15}$$

and

$$\delta \tilde{M}^2 = \begin{pmatrix} \tan \beta & 1 & \frac{\nu_2}{x} \\ 1 & \cot \beta & \frac{\nu_1}{x} \\ \frac{\nu_2}{x} & \frac{\nu_1}{x} & \frac{\nu_1 \nu_2}{x^2} \end{pmatrix} \Delta_p^2. \tag{16}$$

Finally, the charged mass-squared matrix is

$$M_c^2 = \begin{pmatrix} \tan \beta & 1 \\ 1 & \cot \beta \end{pmatrix} (m_4 x - \lambda_7 x^2 - \lambda_4 \nu_1 \nu_2 + \Delta_c^2), \tag{17}$$

where

$$\Delta_c^2 = \frac{3}{16\pi^2} \sum_{m_a \in \{m_{t_1}, m_{t_2}, m_{b_1}\}} m_a^2 \left(\ln \frac{m_a^2}{M_{SUSY}^2} - 1 \right) \frac{\partial^2 m_a^2}{\partial H_1^- \partial H_2^+} \Big|_{vevs}, \tag{18}$$

and

$$\begin{aligned}
\frac{\partial^2 m_{t_1}^2}{\partial H_1^- \partial H_2^+} \Big|_{vevs} &= -\frac{h_t^4 \nu_2^2 (\lambda x)^2 \cot \beta}{(m_{t_1}^2 - m_{t_2}^2)(m_{t_1}^2 - m_{b_1}^2)} - \frac{h_t^2 (\lambda x) A_t}{m_{t_1}^2 - m_{t_2}^2}, \\
\frac{\partial^2 m_{t_2}^2}{\partial H_1^- \partial H_2^+} \Big|_{vevs} &= -\frac{h_t^4 \nu_2^2 (\lambda x)^2 \cot \beta}{(m_{t_2}^2 - m_{b_1}^2)(m_{t_2}^2 - m_{t_1}^2)} + \frac{h_t^2 (\lambda x) A_t}{m_{t_1}^2 - m_{t_2}^2}, \\
\frac{\partial^2 m_{b_1}^2}{\partial H_1^- \partial H_2^+} \Big|_{vevs} &= -\frac{h_t^4 \nu_2^2 (\lambda x)^2 \cot \beta}{(m_{b_1}^2 - m_{t_1}^2)(m_{b_1}^2 - m_{t_2}^2)}.
\end{aligned} \tag{19}$$

The bound on the lightest CP-even Higgs mass is a consequence of the fact that the minimum eigenvalue of M_s^2 is bounded by the minimum eigenvalue of the upper 2×2 submatrix of M_s^2 . Using Eq. (17) to obtain the physical charged Higgs mass squared, m_c^2 , in terms of m_4 ,

$$m_4 x - \lambda_7 x^2 = \frac{1}{2}(m_c^2 + \lambda_4 \nu^2) \sin 2\beta - \Delta_c^2, \tag{20}$$

we may eliminate m_4 from M_s^2 in favour of m_c^2 , and thus write the upper 2×2 submatrix in the form

$$M_s'^2 = M'^2 + \delta M'^2, \tag{21}$$

where

$$M'^2 = \begin{pmatrix} 2\lambda_1 \nu_1^2 & 2(\lambda_3 + \lambda_4) \nu_1 \nu_2 \\ 2(\lambda_3 + \lambda_4) \nu_1 \nu_2 & 2\lambda_2 \nu_2^2 \end{pmatrix} + \begin{pmatrix} \tan \beta & -1 \\ -1 & \cot \beta \end{pmatrix} \frac{1}{2}(m_c^2 + \lambda_4 \nu^2) \sin 2\beta, \tag{22}$$

and

$$\delta M'^2 = \begin{pmatrix} \Delta_{11}^2 & \Delta_{12}^2 \\ \Delta_{12}^2 & \Delta_{22}^2 \end{pmatrix} + \begin{pmatrix} \tan \beta & -1 \\ -1 & \cot \beta \end{pmatrix} (\Delta_p^2 - \Delta_c^2). \tag{23}$$

It is clear from the form of the second term on the right-hand-side of Eq. (23) that the factor $\Delta_p^2 - \Delta_c^2$ does not change the bound at all, but merely serves, for a fixed $\tan \beta$, to shift the bound, when plotted as a function of m_c , to the left or right. We may then drop this term, since its presence may be re-parametrised by a shift in the free parameter m_c . In this way m_{b_1} is eliminated.

From Eq. (22) and Eq. (23) it is a simple matter to determine the shift in the minimum eigenvalue of M'^2 due to squark effects. Defining $A = (M'^2)_{11}$, $B = (M'^2)_{12}$

and $C = (M'^2)_{22}$, the shift is given by

$$\frac{1}{2} \left[\Delta_{11}^2 + \Delta_{22}^2 - \frac{(A - C)(\Delta_{11}^2 - \Delta_{22}^2) + 4B\Delta_{12}^2}{\sqrt{(A - C)^2 + 4B^2}} \right]. \quad (24)$$

Because this shift is a one loop effect, the couplings λ_i in A , B and C may be evaluated at any renormalisation point, since the difference between a coupling evaluated at two different renormalisation points is also a one loop effect, thus giving an overall error at the two loop level, which we neglect in our approximation.

In our previous analysis [7] we used triviality limits on the couplings h_t , λ and k [9] to determine the bound on the lightest CP-even Higgs boson mass m_h^0 not including general squark effects, that is, with $\delta M'^2 \equiv 0$. Let the triviality limit on λ for a given h_t be λ_{max} . In table 1 we reproduce the values of h_t and λ_{max} which generated the bound m_h^0 of our previous analysis — all other values, for a given top quark mass, resulted in smaller values of the bound. It transpires that $k = 0$ in all cases.³ In order to maximise the bound resulting from Eq. (22) we take the limit $m_c = \infty$, thus eliminating m_c as a parameter. As a function of m_c the bound approaches its maximum asymptotically as $m_c \rightarrow \infty$. The approach is rapid, with $m_c \sim 200$ GeV being a good approximation to $m_c = \infty$. Thus, the bound resulting from the purely formal procedure of taking the limit $m_c = \infty$ is not unphysical [7]. Had this not been the case, then certainly we would have an upper bound, but one perhaps not capable of realisation in an actual spectrum.

We use the values of h_t and λ_{max} in table 1 to calculate the shift in the bound on the lightest CP-even Higgs boson mass-squared given by Eq. (24). The shift in the bound on the mass is denoted by δm_h . This is a hideously complicated function of many parameters, including the squark masses m_{t_1} and m_{t_2} , the soft SUSY breaking parameter A_t , the vev of the gauge singlet field x , and m_c . As a function of m_c the

³Taking $k = 0$ gives rise to an axion in the physical spectrum. However, a small, non-zero value of k is sufficient to give the would-be axion a mass of several tens of GeV. Such a value of k does not invalidate our calculations.

Table 1: Lightest CP-even Higgs mass bound in the NMSSM including general squark effects. In row 1 is the top mass, m_t , in GeV; in rows 2 and 3 those values of $h_t(M_{SUSY})$ and $\lambda_{max}(M_{SUSY})$, which produce the bound in our previous analysis, this being in row 4, in GeV; in row 5 is the contribution to the bound resulting from a general squark mass spectrum rather than degenerate squarks with mass M_{SUSY} , in GeV; in row 6 is the lightest CP-even Higgs mass bound including general squark effects, in GeV; in rows 7 and 8 are those value of m_{t_1} and A_t , in GeV, which generate the bound in row 6; $m_{t_2} = 1$ TeV.

m_t	90	100	110	120	130	140	150	160	170	180	190
h_t	0.60	0.67	0.73	0.79	0.84	0.88	0.92	0.95	0.98	1.00	1.03
λ_{max}	0.87	0.85	0.83	0.81	0.79	0.77	0.74	0.71	0.67	0.63	0.50
m_h^0	145	143	140	137	135	132	129	126	124	123	126
δm_h	1	2	3	4	6	8	10	13	16	20	23
m_h	146	145	143	141	140	139	139	139	140	143	149
m_{t_1}	800	770	750	720	700	670	650	630	610	590	570
A_t	940	1000	1040	1130	1160	1270	1330	1410	1500	1630	1820

shift appears, numerically, to be maximised for m_c small, typically between 100 GeV and 200 GeV, with the difference between the shifts at these two points being less than 1 GeV. Thus, we set $m_c = 200$ GeV in the general squark contributions to the bound. Doing this enables us to retain the bound from our previous analysis derived from Eq. (22) in the limit $m_c = \infty$, and simply maximise the squark contributions separately and add them to our previous bound to yield the new bound $m_h = m_h^0 + \delta m_h$. The x dependence is not too strong, with the ratio $r = x/\nu$ typically taking values close to 10 in order to maximise the bound. Where this is not the case, the difference between the bound at its maximum, as a function of x , and that at $r = 10$ is less than 1 GeV. Thus, we set $r = 10$, this giving a reliable indication of the maximum shift. We calculate the shift for various ranges of squark masses and a range of values of A_t , and record the maximum value obtained. The squark masses are allowed to vary between 20 GeV and 1000 GeV in steps of 10 GeV (1000 GeV is the upper limit since we take $M_{SUSY} = 1000$ GeV.) Given this restriction, the value of A_t which maximises the general squark contribution to the bound never exceeds 2 TeV. We impose the constraint $2h_t\nu_2(A_t + \lambda x \cot \beta) \leq |m_{t_1}^2 - m_{t_2}^2|$ which follows from

the form of the squark mass matrix in Eq. (8).

In table 1 and figure 1 we show the bound on the lightest CP-even Higgs mass m_h including general squark effects (and m_h^0 for comparison). Table 1 also shows the values of m_{t_1} and A_t used to generate the bound. As m_t increases, m_{t_1} monotonically decreases, while A_t monotonically increases. The other squark mass, m_{t_2} , takes the value 1 TeV over the whole range of top quark masses — that one squark mass takes on its maximum permitted value to maximise the general squark contribution to the bound can be seen analytically. It will be noticed that these squark masses and A_t are somewhat larger than typically expected from Grand Unification scenarios. However, we feel it wise not to restrict ourselves too excessively to particular prejudices regarding physics as yet unknown (though, of course, the notion of triviality does require the assumption of a SUSY desert up to the unification scale).

Our calculations suggest that the universal upper bound on the lightest CP-even Higgs mass is lifted from 145 GeV to 149 GeV, a mere 4 GeV difference. Table 1 reveals that for larger values of m_t the shifts may be up to 23 GeV, but for small m_t the squark effects are small, as expected. We emphasise that we have not re-maximised the complete one loop corrected bound, but rather made the assumption that the values of h_t and λ_{max} which generated the bound of our previous analysis are not significantly modified by the inclusion of a general spectrum of squark masses.

To conclude, we have supplemented our previous RG analysis of the Higgs sector of the NMSSM with a calculation of general squark effects, and determined how they modify the bound on the mass of the lightest CP-even state. Our calculations indicate that the effects of a general squark spectrum can be very significant for large top quark masses. One reason why, for large top quark masses, squark effects are important is that there exist finite, one-loop diagrams with vertices containing the uncontrolled soft SUSY breaking parameter A_t . These diagrams can give large contributions to

the Higgs boson masses. We close by indicating that in the NMSSM there are similar diagrams involving loops of Higgs bosons, again with vertices containing soft SUSY breaking parameters, this time A_λ and A_k . The existence of these diagrams is directly due to the gauge singlet field N ; they do not occur in the MSSM. We see no reason in principle why these new diagrams should not give rise to similarly large effects, except for obvious factors of 3 due to colour. We are currently in the process of estimating these effects [10].

Acknowledgements

S.F.K. is grateful to the S.E.R.C. for the support of an Advanced Fellowship, and T.E. is grateful to the S.E.R.C. for the support of a studentship.

References

- [1] P. Fayet, Nucl. Phys. **B90** (1975) 104;
J. Ellis, J. Gunion, H. Haber, L. Roszkowski and F. Zwirner, Phys. Rev. **D39** (1989) 844.
- [2] L. Durand and J. Lopez, Phys. Lett. **B217** (1989) 463;
M. Drees, Int. J. Mod. Phys. **A4** (1989) 3635.
- [3] H. Nilles, Phys. Rep. **110** (1984) 1;
H. Haber and G. Kane, Phys. Rep. **117** (1985) 75.
- [4] G. Kane, C. Kolda and J. Wells, Michigan preprint UM-TH-93-24, to appear in Phys. Rev. Lett.;
W. ter Veldhuis, Purdue preprint PURD-TH-92-11;
J. Espinosa and M. Quiros, Phys. Lett. **B302** (1993) 51.

- [5] U. Ellwanger and M. Rausch de Traubenberg, Z. Phys. **C53** (1992) 521;
U. Ellwanger and M. Lindner, Phys. Lett. **B301** (1993) 365.
- [6] U. Ellwanger, Phys. Lett. **B303** (1993) 271.
- [7] T. Elliott, S. King and P. White, Phys. Lett. **B305** (1993) 71.
- [8] J. Ellis, G. Ridolfi and F. Zwirner, Phys. Lett. **B262** (1991) 477;
A. Brignole, J. Ellis, G. Ridolfi and F. Zwirner, Phys. Lett. **B271** (1991) 123.
- [9] P. Binetruy and C. Savoy, Phys. Lett. **B277** (1992) 453;
J. Espinosa and M. Quiros, Phys. Lett. **B279** (1992) 92.
- [10] T. Elliott, S. King and P. White, Southampton preprint SHEP-92/93-21, in preparation.

Figure Caption

Figure 1: The bound on the lightest CP-even Higgs boson mass in the NMSSM. The upper line is the bound m_h including a general spectrum of squark masses. The lower line is the bound m_h^0 of our previous analysis, from assuming degenerate squarks with a common mass M_{SUSY} .